Triggers

Triggers vary from patient to patient, and when possible should be avoided.



ANIMALS

E.g., cats, dogs, horse. Avoid trigger if child/youth is sensitized to an animal and has symptoms around it. If removing the animal from the home is not possible, it should be kept out of the child's bedroom.



COLD AIR

Child/youth can wear a neck warmer or scarf to cover nose and mouth when outdoors.



POLLENS

Staying indoors when pollen counts are high.



E.g., campfire, fireplaces, tobacco/canna-



DUST MITES

Prevention strategies include dust mite mattress and pillow covers.



VIRAL ILLNESSES

The most common trigger. The patient should follow the asthma action plan and take their controller inhaler regularly to prevent the build-up of inflammation. Annual flu shots are recommended.



EXCERCISES

A warm up period before exercise can decrease exercise induced symptoms.

IMPORTANT: Activity should NOT be limited to control symptoms.

There is limited evidence that allergen avoidance and environmental modifications have significant clinical benefit as a treatment strategy for children with asthma. Asthma education should focus on recognizing symptoms, developing an action plan and optimizing adherence to medications.²

Children with asthma often have other allergic conditions (e.g., food allergy, atopic dermatitis, and allergic rhinitis), and suboptimal management of one condition may impact the others.

Helpful Resources

- **<u>QuitNow:</u>** The free quit smoking program of the Government of British Columbia, delivered by British **Columbia Lung Association**
- DrugCocktails.ca: Can help adolescents explore lifestyle choices in private and begin to make their own • health decisions
- **Childhood Asthma:** A Guide for Families and Caregivers: Asthma education video focused on the needs

of Families and Caregivers (easy to remember and share shortened link (bit.ly/pediatric-asthma-video)

- <u>Mandarin</u>
- <u>Cantonese</u>
- <u>Punjabi</u>
- Spanish
- Asthma Wallet card: Patients/families/chosen supports can be provided the Asthma Wallet Card with instructions to bring the card with them the next time they require emergent/urgent care for an asthma exacerbation. The Pediatric Asthma Wallet Care can be completed by the healthcare team before discharge and/or provided to the patient (or family/chosen supports) to complete the information
- First Nations Health Authority Air Purifier Program
 - First Nations Health Authority Air Purifier Support During Wildfire Season Fact Sheet and Contacts
- BC Guidelines. Asthma Diagnosis, Education and Management Table 7. Environmental Controls

References

- 1. Puranik S, Forno E, Bush A, Celedon JC. Predicting severe asthma exacerbations in children. Am J Respir Crit Care Med. 2017;195(7):854-859. DOI: 10.1164/ rccm.201606-1213PP.
- 2. Pedersen SE, Hurd SS, Lemanske RF, et al. Global strategy for the diagnosis and management of asthma in children 5 years and younger. Pediatric Pulmonology. 2011;2010;46:1-17.